

Discrete time survival analysis of longevity in a colony of dog guides

J. B. Cole,^{*}¹ B. R. Southey,² D. E. Franke,³ and E. A. Leighton⁴

¹Animal Improvement Programs Laboratory, Agricultural Research Service,
USDA, Beltsville, MD 20705-2350

jcole@aipl.arsusda.gov

²Department of Animal Sciences, University of Illinois, Urbana-Champaign

³Department of Animal Science, Louisiana State University, Baton Rouge

⁴The Seeing Eye, Inc., Morristown, NJ



Objectives

- ◆ Development of a model for genetic evaluation of working life
- ◆ Variance components estimation for working life
- ◆ Calculation of breeding values

Data

- ◆ Longevity and pedigree data
 - ◆ 954 German shepherds
 - ◆ 1,592 Labrador retrievers
- ◆ All dogs worked as guides
- ◆ Pedigrees complete to founding
- ◆ Breeders selected on an index of hip quality and aptitude

Definition of Working Life

- ◆ Economics favor some measure of total working life
- ◆ Three measures of working life
 - ◆ Graduation to 10 yr (**TWL**)
 - ◆ Graduation to 18 mo (**EWL**)
 - ◆ 18 mo to 10 yr (**LWL**)

Censoring (%) and Working Life (mo)

Trait	GS		LR	
	Censored	WL	Censored	WL
EWL	96.8	56	95.1	67
LWL	49.5	67	44.3	79
TWL	51.1	56	48.9	67

Analysis

- ◆ **Fixed effects**
 - ◆ Duration of time interval (mo or yr)
 - ◆ Contemporary group
 - ◆ Sex of dog
 - ◆ Regression on inbreeding
- ◆ **Random animal effect**
- ◆ **Breeds analyzed separately**

Model

$$\lambda(t_i) = g \left(\sum_{j=1}^k x_{ij} \beta_j \right)$$

$\lambda(\cdot)$ = the discrete hazard function

t_i = the observed time of retirement of individual i

$g(\cdot)$ = link function (logistic or comp log-log)


x_{ij} = the j th explanatory variable for animal i

β_j = the regression coefficient associated with the j th explanatory variable


Survival Models

- ◆ **Logistic link**
 - ◆ Proportional odds model
 - ◆ Variant to duration of the time interval
- ◆ **Complementary log-log link**
 - ◆ Proportional hazards model
 - ◆ Invariant to duration of the time interval
 - ◆ Weibull model when \log_e of duration of the time interval is fit as a covariate

Phenotypic Working Life



Kaplan-Meier Estimated Survival Functions



Variances - Early Working Life (GS)

Model	σ_a^2	σ_p^2	h^2
Proportional odds	0.3	3.6	0.09
Proportional hazards	0.2	1.9	0.12
Weibull	0.5	2.1	0.22

Variances - Early Working Life (LR)

Model	σ_a^2	σ_p^2	h^2
Proportional odds	0.4	3.7	0.12
Proportional hazards	0.3	1.9	0.15
Weibull	1.1	2.7	0.39

Variances - Later Working Life (GS)

Model	σ_a^2	σ_p^2	h^2
Proportional odds	0.6	3.9	0.16
Proportional hazards	0.7	2.3	0.29
Weibull	0.6	2.3	0.28

Variances - Later Working Life (LR)

Model	σ_a^2	σ_p^2	h^2
Proportional odds	1.1	4.4	0.25
Proportional hazards	1.2	2.9	0.42
Weibull	1.2	2.8	0.42

Unequal Use of Parents (GS)

◆ Sires

- ◆ 23 known sires with 767 offspring
- ◆ 3 had > 100 progeny
- ◆ 3 had > 50 progeny
- ◆ 9 had < 20 progeny

◆ Dams

- ◆ 94 known dams
- ◆ 3 had > 20 progeny

- ◆ Inbreeding 0.14 ± 0.001 , 0.0 - 0.38

Unequal Use of Parents (LR)

◆ Sires

- ◆ 41 known sires with 1,187 offspring
- ◆ 5 had > 100 progeny
- ◆ 2 had > 50 progeny
- ◆ 9 had < 20 progeny

◆ Dams

- ◆ 99 known dams
- ◆ 17 had > 20 progeny

◆ Inbreeding 0.07 ± 0.001 , 0.0 - 0.25

Summary

- ◆ The discrete time method produced large estimates of heritability and allowed use of an animal model
- ◆ EBV for early working life may be biased due to censoring
- ◆ Results may reflect founder effects due to the small number of sires